

A summary table of randomised controlled trials comparing
low-carb diets of less than 130g carbohydrate per day
to low-fat diets of less than 35% fat of total calories
with type 2 diabetes participants,
compiled by the Public Health Collaboration

Time Period/Ref.	LC HbA1c Change	LF HbA1c Change	Participants (LC v LF)
1 Year [1]	-8.8 mmol/mol^{^*}	-1.1 mmol/mol	27 v 27
3 Months [2]	-6 mmol/mol[^]	-2.5 mmol/mol	40 v 39
24 Weeks [3]	-16.4 mmol/mol^{^*}	-5.5 mmol/mol	21 v 29
12 Months [4]	-0.2 mmol/mol[^]	+2.6 mmol/mol	47 v 49
24 Months [5]	-1.1 mmol/mol	-2.2 mmol/mol[^]	28 v 40
24 Months [6]	0.0 mmol/mol[^]	+2.2 mmol/mol	30 v 31
24 Weeks [7]	-28.4 mmol/mol^{^*}	-20.8 mmol/mol	46 v 47
6 Months [8]	-4.0 mmol/mol[^]	-1.0 mmol/mol	29 v 30
52 Weeks [9]	-10.9 mmol/mol	-11.0 mmol/mol[^]	41 v 37
6 Months [10]	-7.6 mmol/mol^{^*}	0.0 mmol/mol	30 v 32
4 Months [11]	-9.8 mmol/mol[^]	-4.4 mmol/mol	40 v 36
48 Weeks [12]	-7.7 mmol/mol^{^*}	+1.1 mmol/mol	16 v 21
6 Months [13]	-6.6 mmol/mol^{^*}	-2.2 mmol/mol	12 v 12
3 Months [14]	-6.5 mmol/mol^{^*}	0.0 mmol/mol	16 v 18
32 Weeks [15]	-8.7 mmol/mol^{^*}	-3.3 mmol/mol	12 v 13
12 Weeks [16]	-16.3 mmol/mol^{^*}	0.7 mmol/mol	21 v 12
18 Months [17]	-17.8mmol/mol^{^*}	-11.1mmol/mol	43 v 42
	15/17 LC Are > LF	2/17 LF Are > LC	517 (LC) v 530 (LF)
	10/17 LC Are Sig.	0/17 LF Are Sig.	

[^] = **Greater HbA1c Reduction** * = Significantly Greater HbA1c Reduction Between Groups

References:

[1] The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. May 2004. Stern et al. <https://doi.org/10.7326/0003-4819-140-10-200405180-00007>

[2] Short-term effects of severe dietary carbohydrate-restriction advice in Type 2 diabetes—a randomized controlled trial. September 2005. Daly et al. <https://doi.org/10.1111/j.1464-5491.2005.01760.x>

[3] The effect of a low-carbohydrate, ketogenic diet versus a low-glycemic index diet on glycemic control in type 2 diabetes mellitus. December 2008. Westman et al. <https://doi.org/10.1186/1743-7075-5-36>

[4] Comparative Study of the Effects of a 1-Year Dietary Intervention of a Low-Carbohydrate Diet Versus a Low-Fat Diet on Weight and Glycemic Control in Type 2 Diabetes. July 2009. Davis et al. <https://doi.org/10.2337/dc08-2108>

[5] Effects of a Low-intensity Intervention That Prescribed a Low-carbohydrate vs. a Low-fat Diet in Obese, Diabetic Participants. September 2010. Iqbal et al. <https://doi.org/10.1038/oby.2009.460>

[6] In type 2 diabetes, randomisation to advice to follow a low-carbohydrate diet transiently improves glycaemic control compared with advice to follow a low-fat diet producing a similar weight loss. May 2012. Guldbrand et al. <https://doi.org/10.1007/s00125-012-2567-4>

[7] A very low-carbohydrate, low-saturated fat diet for type 2 diabetes management: a randomized trial. November 2014. Tay et al. <https://doi.org/10.2337/dc14-0845>

[8] Advice to follow a low-carbohydrate diet has a favourable impact on low-grade inflammation in type 2 diabetes compared with advice to follow a low-fat diet. May 2014. Jonasson et al. <https://doi.org/10.3109/07853890.2014.894286>

[9] Comparison of low- and high-carbohydrate diets for type 2 diabetes management: a randomized trial. July 2015. Tay et al. <https://doi.org/10.3945/ajcn.115.112581>

- [10] A randomized controlled trial of 130 g/day low-carbohydrate diet in type 2 diabetes with poor glycaemic control. July 2016. Sato et al. <https://doi.org/10.1016/j.clnu.2016.07.003>
- [11] Short-term safety, tolerability and efficacy of a very low-calorie-ketogenic diet interventional weight loss program versus hypocaloric diet in patients with type 2 diabetes mellitus. September 2016. Goday et al. <https://doi.org/10.1038/nutd.2016.36>
- [12] Two Diets with Different Hemoglobin A1c and Antiglycemic Medication Effects Despite Similar Weight Loss in Type 2 Diabetes. August 2013. Mayer et al. <https://dx.doi.org/10.1111%2Fdom.12191>
- [13] A non-calorie-restricted low-carbohydrate diet is effective as an alternative therapy for patients with type 2 diabetes. 2014. Yamada et al. <https://doi.org/10.2169/internalmedicine.53.0861>
- [14] A Randomized Pilot Trial of a Moderate Carbohydrate Diet Compared to a Very Low Carbohydrate Diet in Overweight or Obese Individuals with Type 2 Diabetes Mellitus or Prediabetes. April 2014. Saslow et al. <https://doi.org/10.1371/journal.pone.0091027>
- [15] An Online Intervention Comparing a Very Low-Carbohydrate Ketogenic Diet and Lifestyle Recommendations Versus a Plate Method Diet in Overweight Individuals With Type 2 Diabetes: A Randomized Controlled Trial. February 2017. Saslow et al. <https://doi.org/10.2196/jmir.5806>
- [16] A food-based, low-energy, low-carbohydrate diet for people with type 2 diabetes in primary care: A randomized controlled feasibility trial. November 2019. <https://doi.org/10.1111/dom.13915>
- [17] Effect of a 90 g/day low-carbohydrate diet on glycaemic control, small, dense low-density lipoprotein and carotid intima-media thickness in type 2 diabetic patients: An 18-month randomised controlled trial. October 2020. <https://doi.org/10.1371/journal.pone.0240158>